

THEORY OF RANDOM MATRICES WITH STRONG LEVEL CONFINEMENT

V. Freilikher[†], E. Kanzieper[†], and I. Yurkevich[‡][†]The Jack and Pearl Resnick Institute of Advanced Technology,
Department of Physics, Bar-Ilan University, Ramat-Gan 52900, Israel[‡]International Centre for Theoretical Physics, Trieste 34100, Italy

(September 11, 1995)

Unitary ensembles of large $n \times n$ random matrices with a non-Gaussian probability distribution $P[\mathbf{H}] \propto \exp\{-\text{tr}V[\mathbf{H}]\}$ are studied using a theory of polynomials orthogonal with respect to exponential weights. Asymptotically exact expressions for density of levels, one- and two-point Green's functions are calculated. We show that in the large- n limit the properly rescaled local eigenvalue correlations are independent of $P[\mathbf{H}]$ while global smoothed connected correlations depend on $P[\mathbf{H}]$ only through the endpoints of spectrum. We also establish previously unknown intimate connection between structure of Szegő function entering strong polynomial asymptotics and mean-field equation by Dyson.

cond-mat/9510002

Random-matrix theory turned out to be general phenomenological approach to description of various phenomena in complex physical systems [1] such as complex nuclei, mesoscopic conductors or quantum chaotic systems. Due to complicated nature of these objects their treatment, based on the direct integration of microscopic equations, runs across considerable obstacles. In this situation the more reasonable way is to deal with the joint distribution function $P[\mathbf{H}]$ of the matrix elements of Hamiltonian $\mathbf{H}$ describing the system involved. If there is not preferential basis in the space of matrix elements $\mathbf{H}_{ij}$ (i.e. the system in question is "as random as possible") one has to require $P[\mathbf{H}]d[\mathbf{H}]$ to be invariant under $\mathcal{R}$ -transformation of the Hamiltonian: $\mathbf{H} \rightarrow \mathcal{R}^{-1}\mathbf{H}\mathcal{R}$. Here $\mathcal{R}$ is orthogonal, unitary or symplectic matrix reflecting the fundamental symmetry of the underlying Hamiltonian. The general form of $P[\mathbf{H}]$ compatible with invariance requirement can be written as $P[\mathbf{H}] = Z^{-1} \exp\{-\text{tr}V[\mathbf{H}]\}$ with $V[\mathbf{H}]$ providing existence of partition function Z . Transformation of $\mathbf{H}$ to diagonal form $\mathbf{H} = \mathcal{S}_\beta^{-1}\mathbf{X}\mathcal{S}_\beta$ and further integration over orthogonal ($\beta = 1$), unitary ($\beta = 2$) or symplectic ($\beta = 4$) group $d\mu(\mathcal{S}_\beta)$ in the construction $P[\mathbf{H}]d[\mathbf{H}]$ yield famous expression for the joint probability density function of the eigenvalues $\{x\}$ of the matrix $\mathbf{H}$:

$$P(\{x\}) = \frac{1}{Z_{n\beta}} \exp \left\{ -\beta \left[\sum_i V(x_i) - \sum_{i < j} \ln |x_i - x_j| \right] \right\}. \quad (1)$$

Confinement potential $V(x)$ contains, in an integral way,

an information about correlations between different matrix elements of a random Hamiltonian $\mathbf{H}$ while logarithmic interaction induced by the invariance requirement is usually treated as a pairwise repulsion between energy levels. We note that in Eq. (1) the symmetry parameter β is factored out from $V(x)$ to fix density of levels in random-matrix ensembles with the same confinement potential but with different underlying symmetry.

The structure of Eq. (1) allows to represent exactly all the global and local statistical characteristics of the physical system with broken time-reversal symmetry ($\beta = 2$) in the terms of polynomials orthogonal with respect to the measure $d\alpha(x) = \exp(-2V(x))dx$.

Hitherto only few cases of strong confinement potentials were *rigorously* treated in the random-matrix theory. Random-matrix ensembles associated with classical orthogonal polynomials have been considered in [2]; those related to some non-classical orthogonal polynomials have been treated in [3], [4]. We also mention the basic work [5] where *ansatz* for functional form of polynomial asymptotics was proposed to describe a class of non-Gaussian unitary random-matrix ensembles.

Since generally an analytical calculation of the corresponding set of orthogonal polynomials is a non-trivial problem, a number of alternative approximate methods were developed. The mean-field approximation proposed by Dyson [6] allows to calculate density of levels in random-matrix ensemble. This approach being combined with the method of functional derivative of Beenakker [7] makes it possible to compute global (smoothed) eigenvalue correlations in large random matrices. Smoothed correlations can also be obtained by diagrammatic approach of Brézin and Zee [8] or by invoking the arguments based on macroscopic electrostatics and linear response theory [9]. We stress that all the methods mentioned above allow to study correlations only in *long-range regime* and, in this sense, they are less informative as compared with the method of orthogonal polynomials [5]. It is worth pointing out the supersymmetry formalism [10], recently developed for invariant random-matrix ensembles with non-Gaussian probability distribution function $P[\mathbf{H}]$, which is exceptional in that it allows to investigate *local* eigenvalue correlations and represents a powerful alternative approach to the classical method of orthogonal polynomials.

It is the aim of this Letter to outline how the problem of non-Gaussian ensembles with unitary symmetry

can be handled by the method of orthogonal polynomials in rather general case. Extended version of the article containing detailed mathematical calculations will be published elsewhere [11]. We stress that our treatment is exact (i.e. it does not involve any conjectures) and based on the recent results [12] obtained in the theory of polynomials orthogonal with respect to exponential weights on $\mathbf{R}$.

Let us consider a class of even confinement potentials $V(x)$ supported on the whole real axis $x \in (-\infty, +\infty)$ which are of *smooth polynomial growth at infinity and increase there at least as $|x|^{1+\delta}$* (δ is arbitrary small positive number). More precisely, we demand $V(x)$ and d^2V/dx^2 be continuous in $x \in (0, +\infty)$, and $dV/dx > 0$ in the same domain of variable x . Further we define the function $T(x) = 1 + x \cdot d^2V/dx^2 \cdot (dV/dx)^{-1}$ and assume that for some $A > 1$ and $B > 1$ the inequality $A \leq T(x) \leq B$ holds for $x \in (0, +\infty)$, and also for x positive and large enough $x^2 \cdot |d^3V/dx^3| \cdot (dV/dx)^{-1} \leq \text{const.}$ The class of potentials $V(x)$ satisfying all the above requirements is said to be of the *Freud type*. The typical examples of the Freud potentials are (i) $V(x) = |x|^\alpha$ with $\alpha > 1$; (ii) $V(x) = |x|^\alpha \ln^\beta(\gamma + x^2)$ with $\alpha > 1$, $\beta \in \mathbf{R}$, and γ large enough.

Defining the Freud (non-Gaussian) measure $d\alpha_{\mathcal{F}}(x) = w_{\mathcal{F}}(x) dx = \exp(-2V(x)) dx$ we can introduce a set of orthogonal polynomials $P_n(x)$

$$\int_{-\infty}^{+\infty} P_n(x) P_m(x) d\alpha_{\mathcal{F}}(x) = \delta_{nm}, \quad (2)$$

for which the following basic result was obtained by D. S. Lubinsky [12]:

$$\lim_{n \rightarrow \infty} \int_{-1}^{+1} d\lambda \left\{ a_n^{1/2} P_n(a_n \lambda) - \left(\frac{2}{\pi} \right)^{1/2} \text{Re} \left[z^n D^{-2} \left(F_n; \frac{1}{z} \right) \right] \right\}^2 w_{\mathcal{F}}(a_n \lambda) = 0. \quad (3)$$

Here parametrizations $z = e^{i\theta}$ and $\lambda = \cos \theta$ were used.

Szegő function $D(g; z)$, appeared in Eq. (3), is of fundamental importance in the whole theory of orthogonal polynomials [13], and takes the form

$$D(g; z) = \exp \left(\frac{1}{4\pi} \int_{-\pi}^{+\pi} d\varphi \frac{1 + ze^{-i\varphi}}{1 - ze^{-i\varphi}} \ln g(\varphi) \right). \quad (4)$$

The first argument of Szegő function in Eq. (3) is

$$F_n(\varphi) = \exp(-V(a_n \cos \varphi)) |\sin \varphi|^{1/2}, \quad (5)$$

and a_n is the n -th Mhaskar-Rahmanov-Saff number being the positive root of the integral equation [14]

$$n = \frac{2a_n}{\pi} \int_0^1 \frac{\lambda d\lambda}{\sqrt{1-\lambda^2}} \left(\frac{dV}{dx} \right)_{x=a_n \lambda}. \quad (6)$$

(In what follows it will be seen that a_n is none other than band edge for eigenvalues of corresponding random-matrix ensemble).

Passing on to the integration over $x = a_n \lambda$ in Eqs. (3) and (6), and leaving aside the problem of remainder term we arrive at the asymptotic formula, $n \gg 1$, for orthogonal polynomials of the Freud type:

$$P_n(x) = \sqrt{\frac{2}{\pi a_n}} \text{Re} \left[z^n D^{-2} \left(F_n; \frac{1}{z} \right) \right], \quad x \in (-a_n, +a_n). \quad (7)$$

Making use of the properties of Szegő function $D(g; e^{i\theta})$ [15] one can obtain the following formula for the orthonormal “wave functions” $\psi_n(x) = P_n(x) \exp(-V(x))$ within the interval $x \in (-a_n, +a_n)$ in the large- n limit:

$$\psi_n(x) = \sqrt{\frac{2}{\pi a_n}} \left[1 - \left(\frac{x}{a_n} \right)^2 \right]^{-1/4} \cos \Phi_n(x), \quad (8)$$

where

$$\Phi_n(x) = \gamma_n(x) + n \arccos \left(\frac{x}{a_n} \right), \quad (9)$$

$$\gamma_n(x) = \frac{1}{2\pi} \mathcal{P} \int_{-a_n}^{+a_n} d\xi \frac{\sqrt{a_n^2 - x^2}}{\sqrt{a_n^2 - \xi^2}} \frac{h(\xi)}{\xi - x}, \quad (10)$$

$$h(\xi) = \frac{1}{2} \ln \left[1 - \left(\frac{\xi}{a_n} \right)^2 \right] - 2V(\xi). \quad (11)$$

In Eq. (10) $\mathcal{P}$ stands for principal value of the integral. Since the same formula takes place for the *Erdős-type* confinement potentials [16] which *grow faster than any polynomial at infinity* (see [17], Ch. 2) the further treatment applies equally to unitary random-matrix ensembles with confinement potentials of the Freud- and Erdős-type. The simple examples of Erdős-type confinement potentials are (i) $V(x) = \exp_k(|x|^\alpha)$ with $\alpha > 0$ and $k \geq 1$ (here $\exp_k$ denotes the exponent iterated k -times); (ii) $V(x) = \exp(\ln^\alpha(\gamma + x^2))$ with $\alpha > 1$, and γ large enough.

All the global and local characteristics of the random-matrix ensembles are determined by two-point kernel [1]

$$K_n(x, y) = \frac{k_{n-1}}{k_n} \frac{\psi_n(y) \psi_{n-1}(x) - \psi_n(x) \psi_{n-1}(y)}{y - x}, \quad (12)$$

where k_n is a leading coefficient of the orthogonal polynomial $P_n(x)$. Substituting Eq. (8) into Eq. (12) we obtain after straightforward calculations in the large- n limit:

$$\begin{aligned}
K_n(x, y) &= \frac{1}{\pi(y-x)} [f_n(x) f_n(y)]^{-1/4} \\
&\times \left\{ \cos \Phi_n(x) \cos \Phi_n(y) \frac{x-y}{a_n} \right. \\
&\quad \left. - \sin \Phi_n(y) \cos \Phi_n(x) \sqrt{f_n(y)} \right. \\
&\quad \left. + \sin \Phi_n(x) \cos \Phi_n(y) \sqrt{f_n(x)} \right\}, \quad (13)
\end{aligned}$$

where $f_n(x) = 1 - (x/a_n)^2$. This equation is valid for arbitrary x and y lying within the band $(-a_n, a_n)$.

Equation (13) allows to compute smoothed (over the rapid oscillations) connected correlations $\nu_c(x, y)$ [5] $\nu_c(x, y) = \langle \nu_n(x) \nu_n(y) \rangle - \langle \nu_n(x) \rangle \langle \nu_n(y) \rangle = -K_n^2(x, y)$, where $x \neq y$ and $\langle \dots \rangle$ denotes averaging over intervals $|\Delta x| \ll a_n$ and $|\Delta y| \ll a_n$ but still containing many eigenlevels. Direct calculations yield the simple universal relationship

$$\nu_c(x, y) = -\frac{1}{2\pi^2} \frac{a_n^2 - xy}{(x-y)^2 \sqrt{a_n^2 - x^2} \sqrt{a_n^2 - y^2}}, \quad x \neq y \quad (14)$$

with dependence on the potential $V(x)$ only through the endpoint a_n of the spectrum.

Now we turn to the local properties of two-point kernel. Assuming that in Eq. (13) $|x-y| \ll a_n$, and both x and y stay away from the band edge a_n we obtain

$$K_n(x, y) = \frac{\sin(\Phi_n(x) - \Phi_n(y))}{\pi(y-x)}. \quad (15)$$

This two-point kernel may be rewritten in locally universal form noting that following representation for $\Phi_n(x)$ takes place

$$\Phi_n(x) = \frac{1}{2} \arccos\left(\frac{x}{a_n}\right) - \pi \int_0^x \omega_{a_n}(\xi) d\xi + \frac{\pi}{4} (2n-1), \quad (16)$$

where

$$\omega_{a_n}(x) = \frac{2}{\pi^2} \mathcal{P} \int_0^{a_n} \frac{\xi d\xi}{\xi^2 - x^2} \frac{dV}{d\xi} \frac{\sqrt{a_n^2 - x^2}}{\sqrt{a_n^2 - \xi^2}}. \quad (17)$$

Since the characteristic scale of the changing of $\omega_{a_n}(\xi)$ is $\varsigma \sim (\omega_{a_n}^{-1} d\omega_{a_n}/d\xi)^{-1} \sim a_n$, Eq. (15) for $|x-y| \ll \varsigma$ is reduced to universal form

$$K_n(x, y) = \frac{\sin[\pi \bar{\nu}_n(y-x)]}{\pi(y-x)} \quad (18)$$

with $\bar{\nu}_n = \omega_{a_n}(\frac{x+y}{2})$ playing the role of local density of levels. Correspondingly, local two-level cluster function being rewritten in rescaled variables s and s'

$$Y_2(s, s') = \left(\frac{K_n^2(x, y)}{\langle \nu_n(x) \rangle \langle \nu_n(y) \rangle} \right)_{\substack{x=x(s) \\ y=y(s')}} = \frac{\sin^2[\pi(s-s')]}{[\pi(s-s')]^2} \quad (19)$$

proves universal Wigner-Dyson level statistics in the unitary random-matrix ensembles with Freud- and Erdős-type confinement potentials (here $s = \bar{\nu}_n x$ and $s' = \bar{\nu}_n y$ are the eigenvalues measured in the local mean level-spacing).

Expression for density of levels $\langle \nu_n(x) \rangle = K_n(x, x)$ immediately follows from Eq. (15), $\langle \nu_n(x) \rangle = -\frac{1}{\pi} d\Phi_n/dx$. It can be written down in two equivalent forms. The first one,

$$\begin{aligned}
\langle \nu_n(x = a_n \cos \theta) \rangle &= \frac{1}{\pi a_n \sin \theta} \\
&\times \frac{d}{d\theta} \left[n\theta + \arg D \left(e^{-2V(a_n \cos \varphi)} |\sin \varphi|; e^{i\theta} \right) \right], \quad (20)
\end{aligned}$$

establishes connection between density of levels in random-matrix ensembles with confinement potentials of Freud- and Erdős-type, and Szegő function, Eq. (4), for corresponding set of orthogonal polynomials.

Another representation of the density of states reads

$$\langle \nu_n(x) \rangle = \frac{2}{\pi^2} \mathcal{P} \int_0^{a_n} \frac{\xi d\xi}{\xi^2 - x^2} \frac{dV}{d\xi} \frac{\sqrt{a_n^2 - x^2}}{\sqrt{a_n^2 - \xi^2}}. \quad (21)$$

Considering this expression as an equation for $V(x)$ one can resolve it invoking the theory of integral equations with Cauchy kernel [18]:

$$V(x) = \int_{-a_n}^{+a_n} dx' \langle \nu_n(x') \rangle \ln|x-x'| + \mu \quad (22)$$

with μ being “chemical potential”. It is non more than famous mean-field equation which, in our treatment, finally follows from the asymptotic formula Eq. (7) for orthogonal polynomials without any physical speculations. Thus, quite surprisingly, Szegő function, Eq. (4), turns out to be closely related to the mean-field approximation by Dyson [6].

Now the calculation of the one-point Green’s function

$$\begin{aligned}
G^p(x) &= \left\langle \text{tr} \frac{1}{x - \mathbf{H} + ip0} \right\rangle \\
&= \mathcal{P} \int_{-a_n}^{+a_n} d\xi \frac{\langle \nu_n(\xi) \rangle}{x - \xi} - i\pi p \langle \nu_n(x) \rangle \quad (23)
\end{aligned}$$

with $p = \pm 1$ becomes quite trivial. We obtain by means of Eqs. (21) and (22):

$$G^p(x) = \frac{dV}{dx} - \frac{2ip}{\pi} \mathcal{P} \int_0^{a_n} \frac{\xi d\xi}{\xi^2 - x^2} \frac{dV}{d\xi} \frac{\sqrt{a_n^2 - x^2}}{\sqrt{a_n^2 - \xi^2}}. \quad (24)$$

Two-point connected Green's function

$$G_c^{pp'}(x, x') = \left\langle \text{tr} \frac{1}{x_p - \mathbf{H}} \text{tr} \frac{1}{x'_{p'} - \mathbf{H}} \right\rangle - \left\langle \text{tr} \frac{1}{x_p - \mathbf{H}} \right\rangle \left\langle \text{tr} \frac{1}{x'_{p'} - \mathbf{H}} \right\rangle, \quad (25)$$

where $x_p = x + ip0$ and $x'_{p'} = x' + ip'0$ ($p, p' = \pm 1$), can also be computed within the proposed formalism. These rather cumbersome calculations [11] lead to the following formula for the smoothed connected two-point Green's function:

$$\overline{G_c^{pp'}}(x, x') = \frac{1}{2} \left\{ pp' \frac{a_n^2 - xx'}{(x - x')^2 \sqrt{a_n^2 - x^2} \sqrt{a_n^2 - x'^2}} - \frac{1}{(x_p - x'_{p'})^2} \right\}, \quad x \neq x'. \quad (26)$$

It also can be shown that the two-point Green's function in the random-matrix theory exhibits the local universality. Introducing normalized and rescaled two-point connected Green's function $g_c^{pp'}(s, s') = G_c^{pp'}(x, x') / \langle \nu_n(x) \rangle \langle \nu_n(x') \rangle$ expressed in the terms of the eigenvalues s and s' measured in the local mean level-spacing, we obtain the following universal relationship that does not depend on the distribution function $P[\mathbf{H}]$:

$$g_c^{pp'}(s, s') = \pi^2 |p - p'| \delta(s - s') + i(p - p') \frac{\sin[\pi(s - s')]}{(s - s')^2} e^{i\pi(s - s') \text{sign}(p - p')}. \quad (27)$$

Note that expression of this type was previously obtained in [19] only for Gaussian random-matrix ensemble using supersymmetry formalism.

In summary, we have presented rigorous analytical consideration of the basic problems of the theory of large Hermitian random matrices with non-Gaussian distribution function $P[\mathbf{H}] \propto \exp\{-\text{tr}V[\mathbf{H}]\}$. Our treatment is applied to very wide class of monotonous confinement potentials which behave *at least* as $|x|^{1+\delta}$ ($\delta > 0$) and *can grow as or even faster than any polynomial at infinity* (note, that this class of confinement potentials is much richer than that considered in [5]). We have shown that in such unitary random-matrix ensembles with non-Gaussian measure $P[\mathbf{H}] d[\mathbf{H}]$ the density of levels and one-point Green's function essentially depend on the measure. In contrast, (connected) two-point characteristics of spectrum are rather universal. Indeed, we have observed global universality of smoothed two-point

connected correlators and local universality of those without smoothing over rapid oscillations. The correlators were shown to depend on the measure only through the endpoints of spectrum (global universality) or through the local density of levels (local universality). Presented rigorous polynomial analysis enabled us to establish new local universal relationship in the random-matrix theory for normalized and rescaled connected two-point Green's function $g_c^{pp'}(s, s')$. It is also worthy of notice an interesting and quite surprising connection between the structure of Szegő function and mean-field equation that has been revealed in the proposed formalism. In our opinion this fact is far from obvious.

One of the authors (E. K.) gratefully acknowledges financial support of The Ministry of Science and The Arts of Israel.

-
- [1] M. L. Mehta, *Random matrices*, (Academic Press, Boston, 1991)
 - [2] T. Nagao and M. Wadati, J. Phys. Soc. Japan **60**, 3298 (1990)
 - [3] L. A. Pastur, Lett. Math. Phys. **25**, 259 (1992)
 - [4] V. Freilikher, E. Kanzieper, and I. Yurkevich, Phys. Rev. E **53**, 2200 (1996)
 - [5] E. Brézin and A. Zee, Nucl. Phys. B[FS] **402**, 613 (1993)
 - [6] F. J. Dyson, J. Math. Phys. **13**, 90 (1972)
 - [7] C. W. J. Beenakker, Phys. Rev. Lett. **70**, 1155 (1993); Phys. Rev. B **47**, 15763 (1993); Nucl. Phys. B[FS] **422**, 515 (1994)
 - [8] E. Brézin and A. Zee, Phys. Rev. E **49**, 2588 (1994)
 - [9] B. Jancovici and P. J. Forrester, Phys. Rev. B **50**, 14599 (1994)
 - [10] G. Hackenbroich and H. A. Weidenmüller, Phys. Rev. Lett. **74**, 4118 (1995)
 - [11] V. Freilikher, E. Kanzieper, and I. Yurkevich (unpublished)
 - [12] D. S. Lubinsky, Acta Appl. Math. **33**, 121 (1993)
 - [13] G. Szegő, Mathematische Annalen **84**, 232 (1921)
 - [14] H. N. Mhaskar and E. B. Saff, Trans. Amer. Math. Soc. **285**, 203 (1984)
 - [15] G. Szegő, *Orthogonal polynomials*, (American Mathematical Society, Providence, 1967), vol. XXIII, Ch. 10
 - [16] Namely, $V(x)$ should be even and continuous in $x \in (-\infty, +\infty)$, d^2V/dx^2 be continuous in $x \in (0, +\infty)$, $dV/dx > 0$ in the same domain of x and continuous at $x = 0$, while function $T(x)$ be positive and increasing in $x \in (0, +\infty)$ with $\lim_{x \rightarrow +0} T(x) > 0$, $\lim_{x \rightarrow \infty} T(x) = \infty$, and $T(x) = \mathcal{O}((dV/dx)^{1/5})$ as $x \rightarrow \infty$. The requirements $(d^2V/dx^2) \cdot (dV/dx)^{-1} \sim V(x)^{-1} \cdot (dV/dx)$ and $|d^3V/dx^3| \cdot (dV/dx)^{-1} \leq \text{const} \cdot (V(x)^{-1} \cdot (dV/dx))^2$ must be fulfilled for x large enough as well.
 - [17] D. S. Lubinsky, *Strong asymptotics for extremal errors and polynomials associated with Erdős-type weights*, Pitman Research Notes in Mathematics **202**, (Longman,

Harlow, Essex, 1989)

- [18] N. I. Akhiezer and I. M. Glazman, *Theory of linear operators in Hilbert space*, (Frederic Ungar Publishing Co., N. Y., 1961), vol. I, Ch. 4
- [19] J. A. Zuk, *Introduction to the Supersymmetry Method for the Gaussian Random-Matrix Ensembles*, `cond-mat/9412060`